




GB 04/00589



INVESTOR IN PEOPLE

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ




I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects company to certain additional company law rules.




*P. Mahoney*

Signed

Dated 9 February 2004

**BEST AVAILABLE COPY**



The  
Patent  
Office

1/77

10 JAN 2003

The Patent Office  
Cardiff Road  
Newport  
Gwent NP9 1RH

**Request for grant of a patent**

(See the notes on the back of this form and also get an explanatory leaflet from the Patent Office to help you fill in this form)

1. Your reference	444.79677/000		
2. Patent application number (The Patent Office will fill in this part)	0300602.0		
3. Full name, address and postcode of the or of each applicant (underline all surnames)	Forinnova AS Thormøhlensgate 55 N-5008 Bergen Norway		
Patents ADP number (if you know it)	8362022001		
If the applicant is a corporate body, give country/state of incorporation	Norway		
4. Title of the invention	COMPOSITIONS		
5. Name of your agent (if you have one)	Frank B. Dehn & Co.		
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	179 Queen Victoria Street London EC4V 4EL		
Patents ADP number (if you know it)	166001 ✓		
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day / month / year)	
8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. See note (d))	Yes		

# Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

0

Description

9

Claim(s)

2

DM

Abstract

-

Drawing(s)

-

10. If you are also filing any of the following, state how many against each item.

Priority documents

-

Translations of priority documents

-

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

-

Request for preliminary examination and search (Patents Form 9/77)

-

Request for substantive examination (Patents Form 10/77)

-

Any other documents (please specify)

-

11.

I/We request the grant of a patent on the basis of this application.

Signature

Frank B Dehn & Co

Date 10 January 2003

12. Name and daytime telephone number of person to contact in the United Kingdom

Julian Cockbain  
020 7206 0600

## Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

## Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s) of the form. Any continuation sheet should be attached to this form.
- If you have answered 'Yes', Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

79677.619

Compositions

5 This invention relates to a composition for use in feeding fish, in particular pre-adult stage fish, especially marine fish larvae and particularly larvae of flatfish, e.g. halibut fry.

10 Marine fish larvae in aquaculture are normally fed with live feed, in particular rotifers (*Bachionus plicatilis*) and *Artemia* (brine shrimp), during the first 4 to 6 weeks of exogenous feeding.

15 *Artemia* for use in aquaculture may be obtained commercially in the dormant egg or cyst form (e.g. from Wudi County Haotian *Artemia* Cysts Co., Ltd of Wudi County, Shandong, China). These may be hatched in aqueous saline (e.g. filtered sea water) and within 18 to 48 hours produce the live baby shrimp or nauplius form which is used as fish feed. The *Artemia* comprise about 48% protein and 18% lipid and thus are generally considered to be an excellent nutrient for fish, including shellfish, in aquaculture.

20 Nonetheless, fish larvae fed on rotifers and *Artemia* have a tendency towards developmental errors in pigmentation and metamorphosis, e.g. incomplete eye migration in flatfish. To some extent this problem has been addressed by enriching rotifer/*Artemia* feeds with marine fish oils (e.g. cod liver oil) prior to distributing the feed to the larvae.

30 The marine fish oils used in this regard are generally triacylglycerols, i.e. triglycerides, and contain a proportion of C<sub>20</sub> and C<sub>22</sub> omega-3 fatty acid (i.e. eicosapentaeneic acid (EPA - C<sub>20</sub>:5n-3) and decosahexaeneic acid (DHA - C<sub>22</sub>:6n-3)) residues.

35 We have now surprisingly found that developmental errors in fish larvae growth can be reduced and survival, growth and feed utilization can be improved by supplementing the live feed with an acylglycerol

composition which contains mono and/or diacylglycerols having C<sub>20</sub> and/or C<sub>22</sub> omega-3 fatty acid residues.

This supplementation may be achieved by allowing the live feed to consume the acylglycerol composition.

5 In effect therefore the live feed then serves in part as a food source for the fish larvae and in part as a carrier for delivering the acylglycerol composition. Simply administering the acylglycerol composition to the fish larvae, e.g. as an emulsion, would result in little  
10 if any being consumed by the fish larvae.

Thus viewed from one aspect the invention provides a fish feed comprising a live feed component which has been fed with an acylglycerol composition which  
15 comprises mono and/or diacylglycerols of at least one fatty acid selected from eicosapentaeneoic acid and docosaheptaeneoic acid, e.g. as at least 5% wt of the total acylglycerol content, especially at least 10% wt, more preferably at least 30% wt, particularly at least  
20 45% wt, for example up to 100% wt.

20 Viewed from a further aspect the invention also provides a process for the preparation of fish feed, said process comprising contacting an acylglycerol composition and a live feed component, characterized in that said acylglycerol composition comprises mono and/or  
25 diacylglycerols of at least one fatty acid selected from eicosapentaeneoic acid and docosaheptaeneoic acid, e.g. as at least 10% wt of the total acylglycerol content, especially at least 20% wt, more preferably at least 30% wt, particularly at least 45% wt, for example up to 100%  
30 wt.

The acylglycerol fed to the live feed component may be a monoacylglycerol, a diacylglycerol, a mixture of mono- and diacylglycerols, a mixture of mono- and triacylglycerols, a mixture of di- and triacylglycerols  
35 or a mixture of mono-, di- and triacylglycerols. Preferably it comprises at least mono and diacylglycerols containing EPA and/or DHA residues, e.g.

containing at least 30% wt, more preferably at least 40% wt, especially at least 50% wt, more especially at least 55% wt of EPA and/or DHA based on the total fatty acid content. Especially preferably at least 25% wt of the EPA, DHA or EPA and DHA residues are present in mono- or diacylglycerols, more especially at least 50% wt. In an especially preferred aspect, the acylglycerol composition also contains acylglycerols having arachidonic acid (AA) residues, preferably in mono and/or diacylglycerols.

Where the EPA or DHA is present in a monoacylglycerol, no other fatty acid residue will be present in the compound. Where however they are present in di- or triacylglycerols, one or two further fatty acid residues will be present. These will preferably be C<sub>16-24</sub> omega-3 or omega-6 acids, especially EPA, DHA or arachidonic acid, i.e. the acylglycerol composition preferably contains diacylglycerol compounds having two EPA or DHA residues, one DHA and one EPA residue, one EPA and one AA residue, or one DHA and one AA residue. Diacylglycerol containing two AA residues and/or monoacylglycerol containing one AA residue may also conveniently be present in the composition. Where the composition also contains triacylglycerols, at least some of these will preferably contain EPA, DHA and/or AA residues.

The DHA content of the acylglycerol composition (relative to total fatty acid content) is preferably at least 10% wt, more preferably at least 20% wt, particularly at least 30% wt, especially at least 45% wt, e.g. up to 100% wt. The EPA content (where EPA is present) is preferably at least 10% wt, more preferably at least 15% wt, e.g. up to 50% wt.

Acylglycerols containing omega-3 acids of marine origin are available commercially, e.g. as EPAX oils from Pronova Biocare of Norway (see [www.pronova.com](http://www.pronova.com)). Examples of available EPAX oils include EPAX 2050 TG

(which has an eicosapentaeneic acid (EPA) content of about 20% wt and a decosahexaeneic acid (DHA) content of about 50% wt and a high mono and diacylglycerol content), EPAX 0626 TG (which has an EPA content of about 6% wt and a DHA content of about 25% wt), EPAX 3000 TG (which has an EPA content of 16-20% wt and a DHA content of 11-13% wt), EPAX 5000 TG (which has an EPA content of about 30% wt and a DHA content of about 20% wt).

10        Acylglycerols containing omega-6 acids are also widely available commercially.

      Mono and diacylglycerols for use according to the invention may be prepared by hydrolysis of omega-3 and/or omega-6 acid containing di and triacylglycerols or by esterification of glycerol or monoacylglycerols with appropriate omega-3 and/or omega-6 fatty acids.

15        Mono- and diacylglycerols containing only DHA, EPA and/or AA residues are themselves novel and form a further aspect of the invention, e.g. in a form substantially free of other acylglycerols, for example at least 90% wt pure, preferably at least 95% wt pure. These can readily be prepared by esterification of glycerol with optionally activated forms of these acids.

20        In the fish feeds of the invention, the live feed component may be any live aquatic animal organism of a size able to be consumed by larval fish, e.g. a monocellular species or a multicellular species having a maximum dimension of up to about 2mm. Preferably the live feed component comprises zooplankton, *Cladocera* (e.g. *Daphnidae*, such as *D. rosea*), rotifers (*Brachionus plicatilis*) or *Artemia*, especially *Artemia*, and in particular *Artemia nauplii*. *Artemia* and other such live feed organisms are available commercially, e.g. in the case of *Artemia* in the dormant cyst form as discussed above. The live feed components will preferably be administered in water, e.g. saline, for example with a salinity (or otherwise expressed osmolality) of from 30

to 120% that of seawater (i.e. 300 to 1200 mOsm/kg).

The live feed compositions according to the invention may conveniently be prepared by addition of an oil-in-water emulsion of the acylglycerols to water containing live feed or by addition of live feed for live feed to an oil-in-water emulsion. The emulsions can be produced by conventional emulsification processes, e.g. by sonification, by using a rotor-stator mixer or by extrusion through a membrane with appropriate pore sizes. Typically droplet sizes from 1 to 100  $\mu\text{m}$ , especially 5 to 50  $\mu\text{m}$  may be used. To assist in emulsion formation, physiologically tolerable emulsification aids, e.g. surfactants such as Tweens, may be added. The aqueous phase of the emulsion will preferably be saline, e.g. as mentioned above.

The acylglycerol will normally be fed to the live feed component at a concentration of about 0.01 to 0.20 g/L, especially 0.05 to 0.10 g/L.

The live feed component will preferably be allowed to consume most or all of the acylglycerol droplets before being fed to the fish or fish larvae. Generally the droplets will be consumed in about 12 to 48 hours, especially about 24 hours, and the live feed component should then be fed to the fish or fish larvae within about 48 hours, preferably within 24 hours, more preferably within 1 hour. If the period between droplet consumption and use as feed is to be prolonged (e.g. over 1 hour), the live feed component is preferably stored under cooling, e.g. at 7 to 8°C. The live feed component will generally be filtered and washed before being fed to the fish or fish larvae.

Besides live feed containing feed compositions, fish at larval and post larval stages may also be fed with formulated feeds containing a mono and/or diacylglycerol composition in accordance with the definitions of the acylglycerol composition given above, e.g. feeds containing further components selected from



proteins, lipids, carbohydrates, colors, vitamins, and minerals.

Thus viewed from a further aspect the invention provides the use in the preparation of a fish feed, e.g. a formulated fish feed, of mono and/or diacylglycerols containing EPA and/or DHA residues.

Viewed from a yet still further aspect the invention provides a method of raising fish comprising feeding fish or fish larvae with an acylglycerol-containing feed, characterized in that said acylglycerol comprises mono and/or diacylglycerols containing EPA and/or DHA residues.

The acylglycerols and/or other lipids may be incorporated into a fish feed in conventional fashion, e.g. by addition to a powdered mixture of other feed components such as for example fish meal, animal and/or vegetable protein, single cell organism protein, cereal flour, gluten, etc, followed by pelletization and drying. Alternatively the acylglycerols and/or other lipids may be combined with minced or emulsified fish, optionally containing other feed components such as animal and/or vegetable protein, single cell organism protein, cereal flour, gluten, etc, extruded, and heated (e.g. with microwave irradiation). The final feed product will typically be in pellet, granule, powder or flake form, preferably granule or powder form, e.g. with a particle size of 50 to 1500  $\mu\text{m}$ , preferably 100 to 1000  $\mu\text{m}$ .

Any variety of fish or fish larvae may be fed according to the invention, including both marine and freshwater fish as well as shellfish and crustaceans, e.g. cod, hake, haddock, halibut, dab, flounder, Japanese flounder, whiting, sole, turbot, sea bass, sea bream, tuna, prawn, shrimp, crab, lobster, crayfish, langoustine, oyster, mussel, scallop, whelk, cockle, etc. The invention however is especially applicable to prawn, shrimp and vertebrate marine fish and

particularly flatfish, i.e. fish which undergo a transformation involving eye migration during their larval stage. The invention is particularly suitable for larval halibut.

5           In the case of filter feeding fish, e.g. shellfish such as bivalves, the acylglycerols may be administered as an emulsion, optionally containing lipid-soluble additives dispersed in the discontinuous oil phase. Thus viewed from a further aspect the invention provides  
10   a method of raising fish comprising feeding filter-feeding fish with an acylglycerol containing feed, characterized in that said feed comprises an oil-in-water emulsion containing mono and/or diacylglycerols of at least one fatty acid selected from EPA and DHA. In  
15   such emulsions, the acylglycerol phase is preferably an acylglycerol or acylglycerol composition as described above and the droplet size is preferably 1 to 100  $\mu\text{m}$ , especially 5 to 50  $\mu\text{m}$ .

          The invention, i.e. the inclusion in feed of mono-  
20   and/or diacylglycerols containing EPA and/or DHA residues is also applicable to feeds for air breathing juvenile vertebrates, e.g. mammals such as sheep, cows, pigs, dogs, cats, humans, etc and birds such as chickens, ducks, geese, turkeys, grouse, pheasants, etc.  
25   In particular it is suitable for use with such juveniles suffering from poor digestive ability. Feed according to the invention for such juveniles will preferably contain the mono- and/or diacylglycerols in the same proportions as described herein for fish feeds,  
30   preferably together with at least one further nutrient selected from carbohydrates, lipids, proteins, protein hydrolysates, vitamins and minerals and mixtures thereof, e.g. milk, fruit, vegetable, meat, fish, cereal, etc.

35           In one particularly preferred embodiment, the mono and/or diacylglycerols will be incorporated into milk powder (e.g. by spray drying milk enriched with the mono

and/or diacylglycerols) or into puréed cooked fruit, vegetable or meat. The quantity of mono and/or diacylglycerols used will typically be 1 to 30% wt, more preferably 2 to 15% wt of the compositions dry weight.

5 Advantageously, such compositions will also contain mono and/or diacylglycerols of C<sub>16-24</sub> omega-6 fatty acids, especially C<sub>18-22</sub> omega-6 fatty acids. Such feed compositions form further aspects of the invention.

10 The invention will now be illustrated by the following non-limiting Example.

Example 1

15 An aqueous emulsion was prepared containing 60% wt. EPAX 2050 TG (available from Pronova Biocare, Norway), 5% wt Tween 20 and 25% wt seawater (33 ppt salt).

20 An equivalent emulsion was prepared using cod liver oil (from Peter Møller AS, Oslo, Norway) in place of the EPAX 2050 TG.

25 *Artemia* cysts were decapsulated and hatched using conventional techniques and at the N\*2 nauplius stage were transferred to enrichment tanks. The emulsions were added to bring the acylglycerol content in the tanks to 0.08 g/L and the tanks were left for 24 hours to allow the artemia to consume the acylglycerols. The acylglycerol enriched *Artemia* were then fed to halibut larvae in triplicate tanks for a period of 77 days.

30 This enriched *Artemia* feed was given to the larvae immediately following the yolk sac stage, i.e. as the first exogenous feed. At 1, 13, 46 and 77 days, larvae were removed and their fatty acid content was measured. At 77 days the mean weight, percentage survival and  
35 percentage eye migration was determined for each group.

The results were as follows:

Acylglycerol		EPAX 2050TG	Cod liver oil
5	Survival (%)	38	11
	Final wet weight (g)	0.24	0.18
	Average eye migration (%)	48	23
	Fatty acid content		
10	(mg/g wet weight) Day 1	7	.7
	Day 13	7	3
	Day 46	41	12
	Day 77	27	13
15	DHA*	8.8	4.4
	EPA*	11.1	10.4
	AA*	3.8	3.7

\* as a percentage of total fatty acid content at 77 days

Claims:

1. A fish feed comprising a live feed component which has been fed with an acylglycerol composition which  
5 comprises mono and/or diacylglycerols of at least one fatty acid selected from eicosapentaeneoic acid and docosahexaeneoic acid.
2. The use in the preparation of a fish feed, e.g. a  
10 formulated fish feed, of mono and/or diacylglycerols containing EPA and/or DHA residues.
3. A method of raising fish comprising feeding fish or fish larvae with an acylglycerol-containing feed,  
15 characterized in that said acylglycerol comprises mono and/or diacylglycerols containing EPA and/or DHA residues.
4. A method of raising fish comprising feeding filter-  
20 feeding fish with an acylglycerol containing feed, characterized in that said feed comprises an oil-in-water emulsion containing mono and/or diacylglycerols of at least one fatty acid selected from EPA and DHA.
- 25 5. An aquatic animal organism having a maximum dimension of 2mm and having in its digestive tracts mono and/or diacylglycerols containing EPA and/or DHA residues.
- 30 6. Milk powder containing mono and/or diacylglycerols containing EPA and/or DHA residues.
7. Puréed cooked fruit, meat or vegetable containing  
35 mono and/or diacylglycerols containing EPA and/or DHA residues.

8. A diacylglycerol wherein both fatty acid groups are EPA, DHA and/or AA residues.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**